





S NCR	of Medicine NLM
Charles Annie Aprile 1991	Nucleotide Protein Genome Structure PopSet Taxonomy OMIM
Scarrein Publik	a/ ⊽ ffor o-me+t-xystky t modification
April 1980-1990 Sept. 1997 Communication and	
	Display Summary Save Text' Order Details Add to Clipboard
endrose PupMrsi	Show: 20 V Items 1-7 of 7 One p
new Pribívlad	1: An H, Wang T, Maier MA, Manoharan M, Ross BS, Cook PD. Related Ar
	Synthesis of novel 3'-c-methylene thymidine and 5-methyluridine/cytidine h-phosphonates and phosphonamidites for new backbone modification of
	oligonucleotides.
O , white	J Org Chem. 2001 Apr 20;66(8):2789-801. PMID: 11304203 [PubMed - in process]
	2: Geary RS, Watanabe TA, Truong L, Freier S, Lesnik EA, Sioufi NB, Sasmor H, Related Ar
	Manoharan M, Levin AA. Pharmacokinetic properties of 2'-O-(2-methoxyethyl)-modified oligonucleotide
	analogs in rats. J Pharmacol Exp Ther. 2001 Mar;296(3):890-7.
	PMID: 11181921 [PubMed - indexed for MEDLINE]
Related Resources	3: Khatsenko O, Morgan R, Truong L, York-Defalco C, Sasmor H, Conklin B, Geary Related Ar
	Absorption of antisense oligonucleotides in rat intestine: effect of chemistry and
	length.
	Antisense Nucleic Acid Drug Dev. 2000 Feb;10(1):35-44. PMID: 10726659 [PubMed - indexed for MEDLINE]
	4: Teplova M, Minasov G, Tereshko V, Inamati GB, Cook Related Articles, Nucleotide, Struc
	PD, Manoharan M, Egli M. Crystal structure and improved antisense properties of 2'-O-(2-methoxyethyl)-RNA
	Nat Struct Biol. 1999 Jun;6(6):535-9. PMID: 10360355 [PubMed - indexed for MEDLINE]
	5: Lind KE, Mohan V, Manoharan M, Ferguson DM. Related Ar
	Structural characteristics of 2'-O-(2-methoxyethyl)-modified nucleic acids from
	molecular dynamics simulations. Nucleic Acids Res. 1998 Aug 15;26(16):3694-799.
	PMID: 9685484 [PubMed - indexed for MEDLINE]
	☐6: Monia BP. Related Ar
	First- and second-generation antisense oligonucleotide inhibitors targeted against
	human c-raf kinase. Ciba Found Symp. 1997;209:107-19; discussion 119-23. Review.
	PMID: 9383572 [PubMed - indexed for MEDLINE]
	7: Bennett GN, Gough GR, Gilham PT. Related Ar

Guanosine tetraphyosphate and its analogues. Chemical synthesis of guanosine 3',5'-dipyrophosphate, deoxyguanosine 3',5'-dipyrophosphate, guanosine 2',5'-bis(methylenediphosphonate), and guanosine 3',5'-bis(methylenediphosphonat Biochemistry. 1976 Oct 19;15(21):4623-8.
PMID: 184820 [PubMed - indexed for MEDLINE]

Display Summary Save Text Order Details Add to Clipboard
Display Summary 1 Gave 16x Section december
Show 20 Items 1-7 of 7
Onow3 1

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Freedom of Information Act | Disclaimer







PubMed	Nucleotide Protein Genome Structure PopSet Taxonomy OMIM
Searca PubMe	d V for single strand mung bean nuclease Got Glean
Eniroz.	Display Summary V Save Text Order Details Add to Clipboard Show: 20" Tems 21-40 of 195 Page 2 of 10 Select page: 1 2 3 4 5 6 7 8
PubMar	21: Sitte N, Saretzki G, von Zglinicki T. Related Ar
NEW 1ªEO√IGO	Accelerated telomere shortening in fibroblasts after extended periods of confluency Free Radic Biol Med. 1998 Apr;24(6):885-93. PMID: 9607597 [PubMed - indexed for MEDLINE]
Schoos	Two superhelix density-dependent DNA transitions detected by changes in DNA adsorption/desorption behavior. Biochemistry. 1998 Apr 7;37(14):4853-62. PMID: 9538002 [PubMed - indexed for MEDLINE]
edition to equest	Preferential accumulation of single-stranded regions in telomeres of human fibrobl Exp Cell Res. 1998 Feb 25;239(1):152-60. PMID: 9511733 [PubMed - indexed for MEDLINE]
	Preferential localization of DNA damage induced by depurination and bleomycin i plasmid containing a scaffold-associated region. Biochem Cell Biol. 1997;75(4):369-75. PMID: 9493959 [PubMed - indexed for MEDLINE]
	The human LON protease binds to mitochondrial promoters in a single-stranded, site-specific, strand-specific manner. Biochemistry. 1998 Feb 17;37(7):1905-9. PMID: 9485316 [PubMed - indexed for MEDLINE]
	26: Yamazaki T, Hasebe T, Shouguchi J, Amano H, Kajiwara S, Shishido K. Structure and function in Escherichia coli of plasmids containing pyrimidine/purine-biased stretch originated from the 5'-flanking region of the basidiomycete ras gene. J Biochem (Tokyo). 1997 Oct;122(4):696-702. PMID: 9399570 [PubMed - indexed for MEDLINE]
	27: Christin L, Wysong DR, Meshulam T, Wang S, Diamond RD. Mechanisms and target sites of damage in killing of Candida albicans hyphae by human polymorphonuclear neutrophils. J Infect Dis. 1997 Dec;176(6):1567-78.

PMID: 9395369 [PubMed - indexed for MEDLINE]

Related Ar **28**: Yu JS, Noll KM. Plasmid pRQ7 from the hyperthermophilic bacterium Thermotoga species strain R replicates by the rolling-circle mechanism. J Bacteriol. 1997 Nov;179(22):7161-4. PMID: 9371465 [PubMed - indexed for MEDLINE] Related Ar 29: Tawa R, Takami M, Imakura Y, Lee KH, Sakurai H. Breaks in double-strand DNA by Cu(II) complexes of etoposide (VP-16) and its derivatives, as evaluated by S1 nuclease treatment. Biol Pharm Bull. 1997 Sep;20(9):1002-5. PMID: 9331985 [PubMed - indexed for MEDLINE] Related Ar 30: Tsurumi T, Yamada H, Daikoku T, Yamashita Y, Nishiyama Y. Strand displacement associated DNA synthesis catalyzed by the Epstein-Barr virus DNA polymerase. Biochem Biophys Res Commun. 1997 Sep 8;238(1):33-8. PMID: 9299446 [PubMed - indexed for MEDLINE] Related Ar 31: Legault J, Tremblay A, Ramotar D, Mirault ME. Clusters of S1 nuclease-hypersensitive sites induced in vivo by DNA damage. Mol Cell Biol. 1997 Sep;17(9):5437-52. PMID: 9271420 [PubMed - indexed for MEDLINE] Related Ar 32: Sutani T, Yanagida M. DNA renaturation activity of the SMC complex implicated in chromosome condensation. Nature. 1997 Aug 21;388(6644):798-801. PMID: 9285594 [PubMed - indexed for MEDLINE] Related Ar 33: Lundin K, Wilichowski E, Ernst BP, Hanefeld F. S1 nuclease hybrid analysis of mitochondrial DNA amplified by long-distance PCI rapid screening for small-scale rearrangements. Nucleic Acids Res. 1997 Jun 15;25(12):2535-6. PMID: 9171111 [PubMed - indexed for MEDLINE] Related Ar 34: Birikh KR, Berlin YA, Soreq H, Eckstein F. Probing accessible sites for ribozymes on human acetylcholinesterase RNA. RNA. 1997 Apr;3(4):429-37. PMID: 9085849 [PubMed - indexed for MEDLINE] Related Ar 35: Carles-Kinch K, Kreuzer KN. RNA-DNA hybrid formation at a bacteriophage T4 replication origin. J Mol Biol. 1997 Mar 14;266(5):915-26. PMID: 9086270 [PubMed - indexed for MEDLINE] Related Ar 36: Furrer P, Bednar J, Stasiak AZ, Katritch V, Michoud D, Stasiak A, Dubochet J. Opposite effect of counterions on the persistence length of nicked and non-nicked DNA. J Mol Biol. 1997 Mar 7;266(4):711-21.

PMID: 9102464 [PubMed - indexed for MEDLINE]

37: Evans E, Fellows J, Coffer A, Wood RD.

Related Ar

Open complex formation around a lesion during nucleotide excision repair provide structure for cleavage by human XPG protein.

EMBO J. 1997 Feb 3;16(3):625-38.

PMID: 9034344 [PubMed - indexed for MEDLINE]

238: Praseuth D, Grigoriev M, Guieysse AL, Pritchard LL, Harel-Bellan A, Nielsen PE, Related Ar Helene C.

Peptide nucleic acids directed to the promoter of the alpha-chain of the interleukin receptor.

Biochim Biophys Acta. 1996 Dec 11;1309(3):226-38. PMID: 8982260 [PubMed - indexed for MEDLINE]

39: Liu B, Maul RS, Kaetzel DM Jr.

Related Articles, Nucleo

Repression of platelet-derived growth factor A-chain gene transcription by an upstr silencer element. Participation by sequence-specific single-stranded DNA-binding proteins.

J Biol Chem. 1996 Oct 18;271(42):26281-90.

PMID: 8824279 [PubMed - indexed for MEDLINE]

40: Bayarsaihan D, Enkhmandakh B, Lukens LN.

Related Ar

Y-box proteins interact with the S1 nuclease-sensitive site in the chicken alpha 2(I) collagen gene promoter.

Biochem J. 1996 Oct 1;319 (Pt 1):203-7.

PMID: 8870670 [PubMed - indexed for MEDLINE]

Display Summary V Save Text Order Details Add to Clipboard Show: 20 V Items 21-40 of 195 Page 2 of 10 Select page: 1 2 3 4 5 6 7 8

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer